Office Use Only				
APPL				
RAD				
CK				

4/11/12



Orthopedic Foundation for Animals
2300 E Nifong Blvd, Columbia, MO 65201-3806
Phone: (573) 442-0418; Fax: (573)875-5073

www.offa.org
A Not-For-Profit Organization

Office Use Only

egistered name:	0	21	Registration number: ♥ AKC □ CKC	Other registry name:
209+ Wai	n Konny		TS 42189201 Date of Birth (month-day-year):	Other registry #:
Lavalie	, ,	sex: Male		
-avalre Number (if any):	_ Tattoo	Deticrochip	9-28-18	
3/27/2/25	4201 017	Concrocnip	Registration number of sire:	Registration number of dam:
JUDGUU TOU UTT			ENCT KO1 12/127	14 ENCI 14131646
Shal	1. 1.			
Owner name:	grove it	125	7-15-19 Examining veterinarian's name or veterinary	hospital:
			East Holm	
iling address:		-	Mailing Address:	23 021 2117
4318 T	R 369		\$5503 C	R 120
n: Ilersbu	State:	H Zip/postal code: 44654	City:	State: Zip/postal code:
one: 330-275	: L mun.		Phone: 330-893-205	E-mail:
(in	aitials of vanistavad	1	imal described on this application to the	public il tile results are abilottila
	ination Results	r),		
			. Classification of luxation	
atellar Exam Normal right	ination Results			kates manually at full extension of the
atellar Exam Normal Fight Patellar Luxatio	ination Results			
atellar Exam Normal right Patellar Luxation bilateral	ination Results		☐ Grade 1—The patella easily lux stifle joint, but returns to the tr☐ Grade 2—There is frequent pa	ochlea when released. tellar luxation which, in some cases
atellar Exam Normal Fight Patellar Luxation bilateral unilateral:	ination Results left on right		☐ Grade 1—The patella easily lux stifle joint, but returns to the tr☐ Grade 2—There is frequent pa becomes more or less permane	ochlea when released. tellar luxation which, in some cases ent.
Normal Night Patellar Luxation bilateral unilateral: luxated:	ination Results eft on right medial	3	☐ Grade 1—The patella easily lux stifle joint, but returns to the tr☐ Grade 2—There is frequent pa becomes more or less permane☐ Grade 3—The patella is perma	ochlea when released. tellar luxation which, in some cases ent. nently luxated with torsion of the tibia
Atellar Exam Normal Fight Patellar Luxation bilateral unilateral: luxated: luxation is:	ination Results left on right medial intermittent	□ left □ lateral □ permanent	☐ Grade 1—The patella easily lux stifle joint, but returns to the tr☐ Grade 2—There is frequent pa becomes more or less permane ☐ Grade 3—The patella is perma and deviation of the tibial crest	ochlea when released. tellar luxation which, in some cases ent. nently luxated with torsion of the tibia t of between 30 degrees and 50 degree
atellar Exam Normal Pight Patellar Luxation bilateral unilateral: luxated:	ination Results left on right medial intermittent <2 months	☐ left ☐ lateral ☐ permanent ☐ 2-6 months	☐ Grade 1—The patella easily lux stifle joint, but returns to the tr ☐ Grade 2—There is frequent pa becomes more or less permane ☐ Grade 3—The patella is perma and deviation of the tibial crest from the cranial/caudal plane.	ochlea when released. tellar luxation which, in some cases ent. nently luxated with torsion of the tibia t of between 30 degrees and 50 degree
Atellar Exam Normal Fight Patellar Luxation bilateral unilateral: luxated: luxation is:	ination Results left on right medial intermittent	□ left □ lateral □ permanent	☐ Grade 1—The patella easily lux stifle joint, but returns to the tr ☐ Grade 2—There is frequent pa becomes more or less permane ☐ Grade 3—The patella is perma and deviation of the tibial crest from the cranial/caudal plane. ☐ Grade 4—The tibia is medially	ochlea when released. tellar luxation which, in some cases ent. nently luxated with torsion of the tibia t of between 30 degrees and 50 degree twisted and the tibial crest may show in the result that it lies 50 degrees to 90
Normal Night Patellar Luxation bilateral unilateral: luxated: luxation is: age of onset:	ination Results left on right left medial left intermittent left con right left on ri	☐ left ☐ lateral ☐ permanent ☐ 2-6 months ☐ > 12 months	☐ Grade 1—The patella easily lux stifle joint, but returns to the treatment of the treatme	ochlea when released. tellar luxation which, in some cases ent. nently luxated with torsion of the tibia t of between 30 degrees and 50 degree twisted and the tibial crest may show in the result that it lies 50 degrees to 90
Normal Night Patellar Luxation bilateral unilateral: luxated: luxation is: age of onset:	ination Results left on right left medial left intermittent left con contact	☐ left ☐ lateral ☐ permanent ☐ 2-6 months ☐ > 12 months	☐ Grade 1—The patella easily lub stifle joint, but returns to the tropy of the common of the common of the tibial cress from the cranial/caudal plane. ☐ Grade 4—The tibia is medially further deviation medially with degrees from the cranial/caudal	ochlea when released. tellar luxation which, in some cases ent. nently luxated with torsion of the tibia t of between 30 degrees and 50 degree twisted and the tibial crest may show in the result that it lies 50 degrees to 90
right Patellar Luxation bilateral unilateral: luxated: luxation is: age of onset: certify that the election of the control of	ination Results left on right medial intermittent <2 months 6-12 months examination was performed on this description.	left lateral permanent 2-6 months > 12 months rmed according to the Coog	☐ Grade 1—The patella easily lux stifle joint, but returns to the treatment of the treatme	tellar luxation which, in some cases ent. nently luxated with torsion of the tibia tof between 30 degrees and 50 degree twisted and the tibial crest may show in the result that it lies 50 degrees to 90 all plane.
Normal Normal Night Patellar Luxation bilateral unilateral: luxated: luxation is: age of onset: Certify that the experimental signs es Animals over A litter of 3 or	ination Results left on right medial intermittent <2 months 6-12 months examination was performed by the complete of the co	left lateral permanent 2-6 months > 12 months rmed according to the Cog I DID NOT veri	Grade 1—The patella easily lunstifle joint, but returns to the transifle joint joint, but returns to the transifle joint joint, but joint joint, but joint joint joint, but joint,	tellar luxation which, in some cases ent. nently luxated with torsion of the tibia tof between 30 degrees and 50 degree twisted and the tibial crest may show in the result that it lies 50 degrees to 90 all plane. group, owned/co-owned by the same person
Normal Normal Night Patellar Luxation bilateral unilateral: luxated: luxation is: age of onset: Certify that the experimental signs es Animals over A litter of 3 or	ination Results left on right medial intermittent <2 months 6-12 months examination was performed by the complete of the co	left lateral permanent 2-6 months > 12 months rmed according to the Cog I DID NOT veri	Grade 1—The patella easily lunstifle joint, but returns to the transifle joint joint, but returns to the transifle joint joint, but joint joint, but joint joint joint, but joint,	tellar luxation which, in some cases ent. nently luxated with torsion of the tibia tof between 30 degrees and 50 degree twisted and the tibial crest may show in the result that it lies 50 degrees to 90 all plane.
Normal Normal Night Patellar Luxation bilateral unilateral: luxated: luxation is: age of onset: Certify that the experimental system A litter of 3 or ms on animals under 1 lible for OFA numbers	ination Results left on right medial intermittent <2 months examination was performed by the complete of th	left lateral permanent 2-6 months > 12 months rmed according to the Cog I DID NOT veri	Grade 1—The patella easily lunstifle joint, but returns to the transifle joint joint, but returns to the transifle joint joint, but joint joint, but joint joint joint, but joint,	cochlea when released. tellar luxation which, in some cases ent. nently luxated with torsion of the tibia tof between 30 degrees and 50 degrees twisted and the tibial crest may show in the result that it lies 50 degrees to 90 all plane. group, owned/co-owned by the same pers